Analysis of the Development of Relativity

Tianyi Jiang

Shanghai Pinghe School, Shanghai, China wildlivfan@gmail.com

Abstract: The advent of relativity theory, initiated by Albert Einstein in the early 1900s, ushered in the era of modern physics by reshaping fundamental concepts of space, time, and gravity. This framework emerged in response to the shortcomings of Newtonian mechanics, particularly its failure to account for the invariance of the speed of light and anomalies in Mercury's orbital path. The paper, through a method of literature review, analyzes and summarizes the historical development and core concepts of both special and general relativity. It explores key background events, major breakthroughs, and their lasting influence on modern science. The paper concludes that the two main contributions of relativity are the famous equations: the mass-equivalence equation in special relativity and the Einstein field equations in general relativity, both of which continue to shape physics today.

Keywords: Relativity, Lorentz Transformation, Mass-energy Equivalence, Spacetime Curvature

1. Introduction

The theory of relativity was prompted by a conflict between classical mechanics and electromagnetic mechanics. Maxwell's equations illustrated that light propagates at a constant speed in a vacuum, which was against the theory of the Ether in classical mechanics. Subsequently, the Michelson-Morley experiment proved the theory of the Ether false. The failure of the theory of ether stimulated the derivation of the Lorentz transformation, which laid the foundation for relativity.

In recent studies, Wolfgang Pauli focused on analyzing how mathematical tools can lead to the derivation of equations in both special and general relativity, while Robert M. Wald focused on specifically general relativity, introducing the derivation of Einstein field equations and the influence of general relativity on cosmology and modern physics.

This paper conducts a literature review of both historical and recent sources to provide an overview of the development of relativity. The paper summarizes key time points of the development of relativity, providing a clear timeline for the readers. In addition, the paper introduces main contributions of relativity, illustrating the significant impact of relativity on modern physics.

2. Background of relativity

There was a long process of preparation before the theory of relativity was proposed. Around the mid-1800s, James Clerk Maxwell introduced a set of equations that bear his name. Expressed in

differential form, these equations are represented as follows:

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\varepsilon_0} \tag{1}$$

$$\nabla \cdot \mathbf{B} = 0 \tag{2}$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial \mathbf{t}} \tag{3}$$

$$\nabla \times \mathbf{B} = \mu_0 \left(\mathbf{J} + \epsilon_0 \frac{\partial \mathbf{B}}{\partial t} \right) \tag{4}$$

with E the electric field, B the magnetic field, ρ the current charge density and J the current density. ϵ_0 is the vacuum permittivity and μ_0 the vacuum permeability.

According to Maxwell's equations, variations in electromagnetic fields—such as those found in light—travel through a vacuum at a fixed speed, denoted by c (299,792,458 m/s) in a vacuum. However, classical mechanics assumed that light propagates through an interstellar medium called the ether, just like sound needs to travel through a medium. If ether did exist, Earth's motion through it should have produced measurable variations in the speed of light.

Conducted in 1887, the Michelson-Morley experiment aimed to identify directional differences in light speed by comparing its propagation along perpendicular paths. The findings, however, revealed no measurable variation, failing to confirm the presence of the hypothesized "ether" [1]. This null result provided compelling evidence against the ether theory and spurred the development of alternative models in theoretical physics.

In 1904, Lorentz proved that Maxwell's equations are invariant under the coordinate transformation:

$$\mathbf{x'} = \kappa \frac{\mathbf{x} - \mathbf{vt}}{\sqrt{(1 - \beta^2)}}, \ \mathbf{y'} = \kappa \mathbf{y}, \ \mathbf{z'} = \kappa \mathbf{z}, \ \mathbf{t'} = \kappa \frac{\mathbf{t} - \left(\frac{\mathbf{v}}{c^2}\right)\mathbf{x}}{\sqrt{1 - \beta^2}} \left(\beta = \frac{\mathbf{v}}{c}\right)$$
(5)

which provided the field intensities in the primed system are suitably chosen in his paper [2]. He concluded the existence of a contraction to all bodies, assuming that translational motion could deform the electrons and that all masses and forces, and purely electromagnetic masses and forces, have the same dependence on the velocity. An imperfection of his theory was that κ must equal to 1. The imperfection was filled by Poincaré. The concept of the Lorentz transformation, which played a significant role in relativity, was first proposed in his paper.

3. Special relativity

In 1905, Albert Einstein introduced the postulates of special relativity in his paper "On the Electrodynamics of Moving Bodies," without referencing earlier works by Lorentz (1904) or Poincaré (1905). The special theory of relativity applies specifically to systems moving at constant velocity relative to one another—in other words, within inertial reference frames [3]. In such frames, no mechanical experiment can reveal which one is "truly" at rest or in motion.

With the introduction of Einstein's relativity principle, the concept of a luminiferous ether lost its credibility. Since it is fundamentally impossible to detect motion relative to the ether, its existence became scientifically irrelevant. Einstein argued that ether should no longer be regarded as a physical medium, but rather as a set of physical properties associated with empty space [4]. The

rejection of the ether hypothesis, along with the notion that space is devoid of matter—essentially a vacuum—led to the conclusion that the speed of light remains constant, regardless of the motion of its source.

The principle of relativity and the constancy of the velocity of light were two basic postulates proposed by Einstein. The two postulates led to the derivation of Lorentz transformation formulae which connect the coordinates (x,y,z,t) in one reference system K to the coordinates (x',y',z',t') in another reference system K', which is moving uniformly relate to K with velocity K' in the positive K' direction. The transformation should be linear because a uniform rectilinear motion one reference system must also be uniform and rectilinear in another reference system. Furthermore, the assumption that the assumption that finite coordinates in K' remain finite in K' supports the validity of Euclidean geometry and the uniformity of space and time. The argument relies on the two postulates, leading to the equation

$$x^2 + y^2 + z^2 - c^2 t^2 = 0 (6)$$

which must also hold in the transformed system

$$x^{2} + y^{2} + z^{2} - c^{2}t^{2} = 0$$
(7)

Since the transformation is linear, it must take the form

$$x^{2} + y^{2} + z^{2} - c^{2}t^{2} = \kappa(x^{2} + y^{2} + z^{2} - c^{2}t^{2})$$
(8)

where κ is a constant that depends on the relative velocity v. On the basis of coordinate transformation, $\kappa=1$, we can deduce Lorentz transformation formulae that

$$x' = \frac{x - vt'}{\sqrt{1 - \beta^2}}, \ y' = y, \ z' = z, \ t' = \frac{t - \left(\frac{v}{c^2}\right)x}{\sqrt{1 - \beta^2}} \left(\beta = \frac{v}{c}\right)$$
 (9)

$$x'^2 + y'^2 + z'^2 - c'^2t'^2 = x^2 + y^2 + z^2 - c^2t^2$$
 (10)

Lorentz transformations explain the relationship between space and time measurements made by two observers in uniform relative motion.

Among the key implications of special relativity are effects such as time dilation and Lorentz contraction. The latter, also referred to as length contraction, describes the observation that an object in motion appears shorter along the direction of motion when measured from a stationary frame, compared to its length when at rest. The contracted length L is given by the formula $L=L_0\sqrt{1-\beta^2}$, where L_0 is the object's proper length. Another consequence of special relativity is time dilation. It's a phenomenon that time dilates for an observer moving relative to another. In other words, time moves more slowly for an observer who is in motion relative to another observer. Analogously, the dilated time t is given by the formula $t=\tau/\sqrt{1-\beta^2}$, where τ is the proper time. At first, time dilation was just a prediction since no experiment could verify its authenticity. Later in 1971, the Hafele-Keating experiment found that the clocks in motion run slower than the stationary clocks at the United States Naval Observatory, which proved the prediction of time dilation.

A major breakthrough of the special theory of relativity is the concept of mass-energy equivalence, which defines the relationship between mass and energy in an object's rest frame. This idea is encapsulated in Einstein's renowned equation $E = mc^2$. It demonstrates that mass can be viewed as a form of energy, meaning that mass and energy are interchangeable and can be represented in either energy or mass units [5]. From the formula $E=mc^2$ we can derive that $\frac{dm}{dt} = \left(\frac{1}{c^2}\right)\left(\frac{dE}{dt}\right)$. The equation implies that even a small amount of mass can correspond to a huge amount of energy. On the contrary, the equation can also be expressed as $\frac{dE}{dt}=\left(c^2\right)\left(\frac{dm}{dt}\right)$, stating that the energy can also contribute to the mass of the system, explaining phenomena in fields such as particle physics and cosmology. In conclusion, the above series of equations illustrates that mass and energy are equivalent and interchangeable, and their units differ by a factor of c^2 , which is a huge number in daily units, causing a huge proportionate change. An important implication of massenergy equivalence is its unification of two fundamental conservation laws in classical physics: the conservation of mass and the conservation of energy. Within the framework of relativity, these are merged into a single law—the conservation of mass-energy. The traditional conservation laws can be seen as limiting cases, valid only when the energy exchanged with the system is minimal compared to its rest mass, making any resulting change in rest mass too small to detect experimentally. This context also gives meaning to the term "special" in special relativity. Einstein regarded his derivation of the mass-energy equivalence formula as the most profound outcome of the special theory of relativity.

4. General relativity

While special relativity applies only to inertial frames, Einstein expanded this framework in 1915 by developing the general theory of relativity, a comprehensive theory of space, time, and gravitation. In his landmark 1916 paper, he introduced the equivalence principle, which asserts that, in a local region, the effects of a uniform gravitational field are indistinguishable from those experienced in an accelerating reference frame. General relativity builds upon and extends special relativity by incorporating non-inertial (accelerated) frames and gravitational phenomena. It reinterprets gravity not as a conventional force, but as the warping of spacetime geometry due to the presence of mass and energy, mathematically described by the metric tensor $g_{\mu\nu}$. The metric tensor $g_{\mu\nu}$ defines the invariant expression for the square of the linear element:

$$ds^2 = g_{\mu\nu} dx_{\mu} dx_{\nu} \tag{11}$$

Within the framework of general relativity, Einstein's field equations establish a connection between the curvature of spacetime and the distribution of matter and energy it contains [6]. These equations are formulated using tensor notation as follows:'

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu} \left(\kappa = \frac{8\pi G}{c^4} \right) \tag{12}$$

In this equation, $G_{\mu\nu}$ denotes the Einstein tensor, which captures the curvature of spacetime; $g_{\mu\nu}$ is the metric tensor that characterizes the geometric structure of spacetime; $T_{\mu\nu}$ represents the stress-energy tensor, which accounts for the distribution of matter and energy; Λ is the cosmological constant; and κ stands for the Einstein gravitational constant. The equation formally

links the local curvature of spacetime (through the Einstein tensor) to the local energy, momentum, and stress (described by the stress-energy tensor).

One of the most significant exact solutions to Einstein's field equations is the Schwarzschild solution, which precisely describes the gravitational field outside a spherically symmetric mass. This solution leads to several deviations from classical Newtonian mechanics, such as the slight anomaly in planetary orbits within our solar system. It also predicts phenomena like the deflection of light by gravity, gravitational redshift, and the time delay of signals passing near massive objects—all of which have been verified through high-precision observations [7].

4.1. Precession of the perihelion of Mercury

Mercury's orbit around the Sun slowly shifts over time, causing the location of its closest approach to the Sun, known as the perihelion, to advance. This gradual rotation of the orbital path is referred to as precession (see Figure 1).




Figure 1. Artist's version of the precession of Mercury's orbit

Observations from Earth show that Mercury's orbit precesses by 5600 arcseconds per century. However, Newtonian mechanics predicts only 5557 arcseconds per century, leaving a discrepancy of 43 arcseconds that classical physics cannot account for. Einstein's general theory of relativity resolves this inconsistency by attributing the additional 43 arcseconds per century to the warping of spacetime near the Sun. Furthermore, the theory also accurately explains a smaller 8.6 arcsecond per century deviation in Venus's perihelion precession, thus addressing another long-standing issue in celestial mechanics.

4.2. Gravitational light bending

Einstein predicted that light passing near a massive object such as the Sun would bend due to spacetime curvature. Light passing near a mass M is bent by

$$\Delta \phi = \frac{4GM}{c^2b} \tag{13}$$

where b is the closest approach distance. Einstein calculated that the deflection of light passing close to the sun predicted by his theory would be 1.75 arcseconds, twice the Newtonian value, which was later confirmed by Arthur Eddington's solar eclipse observations in 1919.

4.3. Gravitational redshift

According to Einstein's general relativity, electromagnetic waves experience an increase in wavelength, known as gravitational redshift, when they move away from a gravitational field [8]. This phenomenon occurs because photons lose energy while escaping a gravitational potential, even though their speed remains unchanged. Conversely, when photons move deeper into a gravitational field, they gain energy, resulting in what is called gravitational blueshift. The gravitational redshift effect has been experimentally confirmed, most notably by the Pound-Rebka experiment, which precisely measured the redshift of photons as they ascended out of Earth's gravitational field.

4.4. Gravitational time delay

Gravitational time delay, commonly referred to as the Shapiro delay, describes the additional time taken by light as it passes near a massive body, a result of time dilation predicted by general relativity [9]. Specifically, radar signals that travel close to a massive object require slightly more time to return than they would in flat spacetime. This phenomenon was first confirmed by Irwin Shapiro through measurements of radar echoes reflected from Venus and Mercury.

In conclusion, studying the curvature of spacetime and null geodesics within the Schwarzschild geometry yields several key predictions: the advance of planetary perihelia, the deflection of light by gravity, gravitational redshift, and the time delay of radar waves. Each of these effects has been empirically validated through observations within our solar system, providing strong support for the accuracy of general relativity.

5. Conclusion

This research contributes to the existing literature by marking key time points and theoretical contributions in the development of relativity. This study has practical significance for the field of physics education by providing a clear timeline of the development of relativity. The findings hold educational value, especially for newcomers of physics seeking an accessible entry point into complex physical theories. Nonetheless, this study lacks the derivation of equations and details at each stage of development, which may affect the generalizability of the findings. In the future, the author will investigate the basis of each equation and how the theory of relativity can relate to current technology. Overall, this study provides a concise yet informative overview of relativity's evolution and underscores the importance of multi-source literature review in constructing a comprehensive understanding of scientific progress.

References

- [1] Boeglin, W. (2025). Modern Lab Experiments: 16. Michelson Interferometer wanda.fiu.edu/boeglinw/courses/Modern lab manual3/michelson.html.
- [2] Pauli, W. (2013). Theory of relativity. Courier Corporation.

Proceedings of CONF-APMM 2025 Symposium: Simulation and Theory of Differential-Integral Equation in Applied Physics DOI: 10.54254/2753-8818/2025.DL25567

- [3] The Editors of Encyclopaedia Britannica. (2010). Special relativity | Definition & Equation. Encyclopedia Britannica. https://www.britannica.com/science/special-relativity
- [4] Einstein, A., & Einstein, A. (1920). Äther und Relativitäts-theorie (pp. 1-15). Springer Berlin Heidelberg.
- [5] Resnick, R. (1991). Introduction to special relativity. John Wiley & Sons.
- [6] Einstein, A., Perrett, W., and Jeffery, G. (1916) The foundation of the general theory of relativity. Annalen der Physik 354.7: 769.
- [7] Wald, R. M. (2010). General relativity. University of Chicago press.
- [8] Cosmos. (n.d.). Gravitational Redshift. https://astronomy.swin.edu.au/cosmos/g/Gravitational+Redshift
- [9] Cosmos. (n.d.). Shapiro Delay. https://astronomy.swin.edu.au/cosmos/s/Shapiro+Delay